

Markscheme

November 2025

Chemistry

Higher level

Paper 2

© International Baccalaureate Organization 2025

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2025

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2025

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.



Subject Details: Chemistry Higher Level Paper 2 Markscheme

Candidates are required to answer **ALL** questions. Maximum total = **[90 marks]**.


1. Each row in the “Question” column relates to the smallest subpart of the question.
2. The maximum mark for each question subpart is indicated in the “Total” column.
3. Each marking point in the “Answers” column is shown by means of a tick (✓) at the end of the marking point.
4. A question subpart may have more marking points than the total allows. This will be indicated by “**max**” written after the mark in the “Total” column. The related rubric, if necessary, will be outlined in the “Notes” column.
5. An alternative word is indicated in the “Answers” column by a slash (/). Either word can be accepted.
6. An alternative answer is indicated in the “Answers” column by “**OR**”. Either answer can be accepted.
7. An alternative markscheme is indicated in the “Answers” column under heading **ALTERNATIVE 1** etc. Either alternative can be accepted.
8. Words inside chevrons « » in the “Answers” column are not necessary to gain the mark.
9. Words that are underlined are essential for the mark.
10. The order of marking points does not have to be as in the “Answers” column, unless stated otherwise in the “Notes” column.
11. If the candidate’s answer has the same “meaning” or can be clearly interpreted as being of equivalent significance, detail and validity as that in the “Answers” column then award the mark. Where this point is considered to be particularly relevant in a question it is emphasized by **OWTTE** (or words to that effect) in the “Notes” column.
12. Remember that many candidates are writing in a second language. Effective communication is more important than grammatical accuracy.
13. Occasionally, a part of a question may require an answer that is required for subsequent marking points. If an error is made in the first marking point then it should be penalized. However, if the incorrect answer is used correctly in subsequent marking points then **follow through** marks should be awarded. When marking, indicate this by adding **ECF** (error carried forward) on the script.
14. Do **not** penalize candidates for errors in units or significant figures, **unless** it is specifically referred to in the “Notes” column.
15. If a question specifically asks for the name of a substance, do not award a mark for a correct formula unless directed otherwise in the “Notes” column. Similarly, if the formula is specifically asked for, do not award a mark for a correct name unless directed otherwise in the “Notes” column.
16. If a question asks for an equation for a reaction, a balanced symbol equation is usually expected, do not award a mark for a word equation or an unbalanced equation unless directed otherwise in the “Notes” column.



Ignore missing or incorrect state symbols in an equation unless directed otherwise in the “Notes” column.


Question			Answers	Notes	Total
1.	(a)		Any one of: forms nitrous acid / HNO_2 / nitric acid / HNO_3 ✓ contributes to acid rain ✓ catalyses decomposition of ozone / contributes to ozone hole/depletion ✓ «respiratory» irritant/toxic ✓	Accept suitable equations showing acid rain formation or ozone depletion. Accept photochemical smog/forms NO_2.	1 max
1.	(b)		$T = 200 + 273$ «K» ✓ «$n = PV/RT$» «$n = \frac{(1.0 \times 10^5)(1.0 \times 10^{-3})}{8.31(200+273)}$» «$n =$» 0.025 «mol» ✓ «$0.10\% \times 0.025 =$» 2.5×10^{-5} «mol» ✓	Award [3] for correct final answer.	3
1.	(c)		NO polar AND N_2 non-polar OR NO has dipole-dipole «and LDF» AND N_2 has London/dispersion «forces between molecules» ✓ stronger intermolecular forces in NO OR dipole-dipole «forces» are stronger «than LDF» ✓		2
1.	(d)	(i)	Initial: +2 AND Final: 0 ✓	Accept II for initial oxidation state, but do not accept 2+ or 2.	1
1.	(d)	(ii)	$2\text{NO(g)} + 2\text{CO(g)} \rightarrow 2\text{CO}_2\text{(g)} + \text{N}_2\text{(g)}$ ✓		1

Question			Answers	Notes	Total
2.	(a)	(i)		Accept other forms of labelling instead of an arrow. Mark is for identifying the bond.	1
2.	(a)	(ii)	$\text{Na}_2\text{CO}_3(\text{s}) + 2\text{HNO}_3(\text{aq}) \rightarrow 2\text{NaNO}_3(\text{aq}) + \text{H}_2\text{O}(\text{l}) + \text{CO}_2(\text{g})$ ✓		1
2.	(b)	(i)		Accept any combination of dots or crosses to represent electrons, or lines to represent electron pairs. Charge must be shown, but square brackets are not required. Do not accept delocalized structure.	1
2.	(b)	(ii)	$\text{Ni}(\text{NO}_3)_2$ ✓		1
2.	(b)	(iii)	resonance / delocalization «of π -electrons» ✓		1
2.	(b)	(iv)	124 «pm» ✓	Accept range $114 < x < 136$ «pm».	1


2.	(c)		<table><tr><td>Metal added:</td><td colspan="2">Mn</td><td>Cu</td></tr><tr><td>Observations:</td><td>«solution» colour changes OR solid/metal product / coat «on metal»</td><td>AND</td><td>no change «in solution colour» OR no solid/metal produced / no coat «on metal» ✓</td></tr></table> Explanation: Mn lower/more negative $E^\ominus$ AND Cu higher/more positive $E^\ominus$ than Ni OR Mn more reactive/readily oxidized AND Cu less reactive/readily oxidized than Ni OR Mn stronger reducing agent AND Cu weaker reducing agent than Ni ✓	Metal added:	Mn		Cu	Observations:	«solution» colour changes OR solid/metal product / coat «on metal»	AND	no change «in solution colour» OR no solid/metal produced / no coat «on metal» ✓	Accept “no reaction” for Cu observation.	Award [1 max] for correct observation and explanation for one metal only.	2
Metal added:	Mn		Cu											
Observations:	«solution» colour changes OR solid/metal product / coat «on metal»	AND	no change «in solution colour» OR no solid/metal produced / no coat «on metal» ✓											
2.	(d)	(i)	Lewis formula 1: O –1 AND P +1 «are the only formal charges» ✓ Lewis formula 2: formal charges zero/closest to zero «and hence preferred» ✓	Accept labelling for M1: $\begin{array}{c} \text{:}\ddot{\text{O}}\text{:}^\ominus \\ \\ \text{H}-\ddot{\text{O}}-\text{P}^\oplus-\ddot{\text{O}}-\text{H} \\ \\ \text{:}\ddot{\text{O}}\text{:} \\ \\ \text{H} \end{array}$		2								
2.	(d)	(ii)	H ₂ PO ₄ [–] ✓			1								

Question			Answers	Notes	Total
3.	(a)	(i)	«mol styrene = $\frac{90}{100} \times \frac{1000}{8(12.01)+10(1.01)}$ » «mol styrene =» 8.48 «mol» ✓ «mass styrene = $8.48 \times (8(12.01) + 8(1.01)) = 882.88$» «mass styrene =» 880 «g» ✓	Award [2] for correct final answer in range 880-890 «g».	2
3.	(a)	(ii)	«$100 \times \frac{104.16}{106.18}$ » = 98.1% ✓		1
3.	(b)		 two curves, each passing through a maximum AND reaching same energy level ✓ endothermic enthalpy change labelled ✓ both activation energies correctly labelled ✓	Do not penalize curve showing multiple steps for the catalysis in M1. Accept double-headed arrows or lines in M2 and M3, but not arrows pointing down.	3

3.	(c)	(i)	 Proportion of particles Kinetic energy 900 K curve with peak higher AND to left AND decreasing at lower KE ✓	<i>Do not award mark if line crosses x axis or rises above the 900K curve at high energy.</i>	1
3.	(c)	(ii)	 Proportion of particles Kinetic energy 900 K E_a ✓	<i>Accept a correct label on x-axis or on a drawn line.</i>	1
3.	(c)	(iii)	correct reference to area under curve «to right of E_a» ✓ fewer particles with $E \geq E_a$ ✓		2


3.	(c)	(iv)	shifts left/to reactants AND more moles/molecules «gas» in products ✓		1
3.	(c)	(v)	«forward» reaction is endothermic AND shifts to left/reactants OR $E_{a \text{ fwd}} > E_{a \text{ rev}}$, so forward reaction more affected by temperature AND shifts to left/reactants ✓	Accept “exothermic reverse reaction is favoured”.	1
3.	(d)	(i)	 straight line through origin ✓		1
3.	(d)	(ii)	<i>Rate equation:</i> «rate =» $k [\text{C}_8\text{H}_{10}]$ ✓ <i>Units of k:</i> s^{-1} ✓		2

3.	(f)	(i)	$\frac{[C_6H_5COO^-][H^+]}{[C_6H_5COOH]}$	Accept H_3O^+ for H^+ .	1
3.	(f)	(ii)	horizontal line from point on curve at 10 cm ³ of NaOH to y-axis ✓	Accept line intersecting y-axis between 4.1 and 4.4.	1
3.	(f)	(iii)	bromothymol blue OR phenol red OR phenolphthalein ✓		1
3.	(f)	(iv)	«benzoate ion» hydrolysis/reaction with water AND forms OH ⁻ OR $C_6H_5COO^-(aq) + H_2O(l) \rightleftharpoons C_6H_5COOH(aq) + OH^-(aq)$ ✓		1
3.	(f)	(v)	«(0.020 × 0.010) ÷ 0.010 =» 0.020 «mol dm ⁻³ » ✓		1
3.	(g)	(i)	partially neutralize benzoic acid with NaOH OR amount/moles of NaOH added is smaller «than amount of benzoic acid» ✓		1
3.	(g)	(ii)	remains constant «as [A ⁻] : [HA] ratio is unchanged» ✓	Do not accept “slight/minimal change”.	1

3.	(h)	(i)	condensation OR polyester ✓		1
3.	(h)	(ii)	 ester linkage ✓ continuation bonds AND rest of structure ✓	Accept without square brackets or <i>n</i>. Do not award M2 if more than one repeating unit is included.	2

Question			Answers	Notes	Total
4.	(a)	(i)	electrophilic addition ✓		1
4.	(a)	(ii)	high electron density of C=C/pi bond ✓	Do not accept only “presence of C=C”.	1

4.	(b)	(i)	ALTERNATIVE 1: «bonds broken» $\text{C}=\text{C} + \text{H}-\text{Br} / 614 + 366 / 980 \text{ «kJ mol}^{-1}\text{»}$ AND «bonds formed» $\text{C}-\text{C} + \text{C}-\text{H} + \text{C}-\text{Br} / 346 + 414 + 285 / 1045 \text{ «kJ mol}^{-1}\text{»} \checkmark$ «$\Delta H = 980 - 1045 =$» $-65 \text{ «kJ mol}^{-1}\text{»} \checkmark$ ALTERNATIVE 2: «bonds broken» $1\text{C}=\text{C} + 4\text{C}-\text{H} + 1\text{H}-\text{Br} / 614 + 4 \times 414 + 366 / 2636 \text{ «kJ mol}^{-1}\text{»}$ AND «bonds formed» $1\text{C}-\text{C} + 5\text{C}-\text{H} + 1\text{C}-\text{Br} / 346 + 5 \times 414 + 285 / 2701 \text{ «kJ mol}^{-1}\text{»} \checkmark$ «$\Delta H = 2636 - 2701 =$» $-65 \text{ «kJ mol}^{-1}\text{»} \checkmark$	<i>Award [2] for correct final answer.</i> <i>Award [1 max] for $+65 \text{ «kJ mol}^{-1}\text{»}$.</i>	2
4.	(b)	(ii)	Any two of: bond enthalpies are average values $\checkmark$ enthalpies of formation are specific to the compounds $\checkmark$ bond enthalpies apply to gases AND this reaction involves a liquid $\checkmark$ bond enthalpies do not account for the change of state $\checkmark$		2 max
4.	(b)	(iii)	two moles/molecules of reactants form one mole/molecule of product OR gaseous reactants form liquid product $\checkmark$		1

4.	(b)	(iv)	$\Delta S = 199 - 220 - 199 = -220 \text{ J K}^{-1} \text{ mol}^{-1}$ ✓ $\Delta G^\ominus = -65\,000 \text{ J mol}^{-1} - (298 \text{ K} \times -220 \text{ J K}^{-1} \text{ mol}^{-1})$ » $\Delta G^\ominus = +560 \text{ J mol}^{-1} / +0.56 \text{ kJ mol}^{-1}$ AND non-spontaneous ✓	Award [2] for correct final answer ($\Delta G^\ominus$ value and spontaneity). If $\Delta H = -58 \text{ kJ mol}^{-1}$ used, answer is «+»7600 «J mol⁻¹» / «+»7.6 «kJ mol⁻¹».	2
4.	(c)	(i)	nucleophilic substitution ✓	Accept S_N2 , but not S_N1 .	1
4.	(c)	(ii)	 curly arrow from lone pair/negative charge on O in HO^- to C ✓ curly arrow from C–Br bond to Br ✓ transition state showing negative charge AND partial bonds ✓	Award [2 max] for correct S_N1 mechanism if not stated in (i). Accept curly arrow going from C–Br bond to Br either in bromoethane or in the transition state for M2. Do not penalize if HO and Br are not at 180°, or the square bracket is missing, in M3. Do not award M3 if OH ---- C connection is seen.	3
4.	(c)	(iii)	primary halogenoalkane ✓	Accept reasons why, such as lack of steric hindrance or less stable carbocation.	1
4.	(c)	(iv)	2 / second ✓		1

4.	(c)	(v)	iodoethane AND «C–I» bond weaker/easier to break ✓	Accept “iodoethane AND I is a better leaving group”.	1
4.	(c)	(vi)	quartet / quadruplet / 4 ✓		1
4.	(c)	(vii)	bromoethane AND absence of O–H «peak» AND in 3200–3600 «cm ⁻¹ range» OR bromoethane AND C–Br «peak» AND 500–600 «cm ⁻¹ » ✓	Accept specific value in 500 – 600 «cm ⁻¹ » range.	1

Question			Answers	Notes	Total
5.	(a)		«two» more neutrons/n in ⁸¹ Br ✓		1
5.	(b)	(i)	Any two of: «Br ₂ molecules made up of» ⁷⁹ Br – ⁷⁹ Br «158 peak» AND ⁷⁹ Br – ⁸¹ Br «160 peak» AND ⁸¹ Br – ⁸¹ Br «162 peak» ✓ number/probability/abundance of ⁷⁹ Br – ⁷⁹ Br and ⁸¹ Br – ⁸¹ Br «molecules» is «nearly» same «hence peaks same height» ✓ number/probability/abundance of ⁷⁹ Br – ⁸¹ Br «molecules» is double «of ⁷⁹ Br – ⁷⁹ Br and ⁸¹ Br – ⁸¹ Br molecules hence peak is double the height» ✓		2 max

5.	(b)	(ii)	$79 \times 50.75\% + 81 \times 49.25\% \checkmark$ $= 79.99 \checkmark$	<i>Award [2] for correct final answer.</i> <i>Do not accept 79.90, the value in the data booklet.</i>	2
5.	(c)	(i)	«electronegativity difference $\Rightarrow$ 2.0 AND «average electronegativity $\Rightarrow$ 2.0 $\checkmark$ ionic AND 35% covalent $\checkmark$	<i>Accept range 30 - 40% for percentage covalent character.</i>	2

5.	(c)	(ii)	three values from data booklet: «Br–Br enthalpy» 193 AND «1st IE Ca» 590 AND «EA of Br» –325 «kJ mol^{–1}» ✓ –325 x 2 / –650 «kJ mol^{–1}» ✓ 648 + 178 + 31 + 193 + 590 + 1145 = ΔH_{latt.} + 650 AND «ΔH_{latt.} = +2135 = +»2100 «kJ mol^{–1}» ✓	<i>Award [3] for correct final answer.</i> <i>Award [2 max] for –2100 «kJ mol^{–1}».</i> <i>M1 and M2 can be scored on the diagram.</i> <i>Award ECF for M3 for correct application of Hess's law based on the candidate's values/cycle.</i>	3
----	-----	------	---	--	---

5.	(c)	(iii)	Ca^{2+} and Co^{2+} lost $4s^2$ electrons «so smaller than Br^-» OR Ca^{2+} and Co^{2+} have electrons in 3 shells/energy levels «so smaller than Br^-» ✓ Co^{2+} has greater «effective» nuclear charge/more protons than Ca^{2+} «so smaller» ✓ Br^- has electrons in 4 shells/energy levels / $4p^6$ «so larger than cations» ✓		3
5.	(c)	(iv)	cobalt(II) bromide AND «cation has» higher charge density / smaller radius ✓		1
5.	(d)	(i)	$[\text{Ar}] 4s^2 3d^7$ / $[\text{Ar}] 3d^7 4s^2$ ✓		1
5.	(d)	(ii)	incomplete/partially-filled d «sub-level» ✓		1
5.	(d)	(iii)	«wavelength = $3.00 \times 10^8 \text{ ms}^{-1} \div 4.5 \times 10^{14} \text{ s}^{-1} \Rightarrow 6.7 \times 10^{-7} \text{ m}$» / 670 «nm» ✓ red is absorbed AND green is complementary ✓	Accept any units for wavelength in M1.	2

5.	(e)	(i)	<i>Anode:</i> $\text{Co(s)} \rightarrow \text{Co}^{2+}(\text{aq}) + 2\text{e}^-$ ✓ <i>Cathode:</i> $\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co(s)}$ ✓	<i>Award [1 max] for correct half-equations at incorrect electrodes.</i>	2
5.	(e)	(ii)	$\text{Cl}_2(\text{g}) + 2\text{Br}^-(\text{aq}) \rightarrow 2\text{Cl}^-(\text{aq}) + \text{Br}_2(\text{aq})$ ✓		1
